




Bi/Tri/Quad MIX INTRACAVERNOSAL

SELF-INJECTION GUIDE

Patient Guide

YOUR PRESCRIPTION: Inject the prescribed dose amount into the penis as directed by your prescriber.	
My usual injection time: _____	Dose/strength: See pharmacy label

1. Prepare  - Remove medication vial from freezer. Allow medication to thaw. - Wash and dry your hands. - Gather your medication, syringe, alcohol pad, and gauze.	2. Check the Medication  - Confirm your name, medicine, dose/strength, and beyond-use date. - Do not use it if cloudy, discolored, leaking, or if the solution contains particles. Call the pharmacy.	3. Draw up Medication  - Remove the cap from vial and syringe. - Wipe rubber stopper of vial with alcohol wipe. Let dry. - Insert syringe and invert vial. - Draw medication up by pulling down on the plunger on syringe. Stop when you reach the prescribed dose amount.
4. Choose a site  - Inject into the side of the shaft, close to the base of the penis at the two or ten o'clock position. (avoiding the top and bottom). - Avoid veins and the head (glans). - Be sure to rotate injection sites.	5. Clean and inject  - Clean the injection site with an alcohol pad and let it dry. - Hold the penis steady and insert the needle at a 90° angle (straight) into the side of the shaft. - Avoid twisting the penis.	6. After the injection  - Remove the needle straight out. - Apply gentle pressure with gauze for about 1 minute. Do not rub. - Dispose of needle in a sharps container or closed plastic container

IMPORTANT SAFETY INFORMATION

Read this page before your first dose and keep it with your medication.

⚠ POSSIBLE SIDE EFFECTS - Headache, facial flushing, nasal congestion, upset stomach, dizziness. - Penile pain, prolonged erection (priapism), bleeding, or bruising at the injection site. - Contact your physician if you experience redness, lumps, swelling, tenderness, or curving of the penis or a prolonged painful erection (priapism).	+ GET EMERGENCY HELP NOW - If you have an erection that lasts more than 4 hours. - Priapism is a medical emergency and can cause permanent damage.	☎ CALL YOUR PHYSICIAN PROMPTLY - If you experience chest pain, shortness of breath, or irregular heartbeat. - If you have sudden vision loss. - If you have pain or discomfort that does not improve. - If you notice any changes in your condition.
❄ STORAGE - Store in the freezer until ready for use. - Keep in the original box to protect from light. - Keep out of reach of children. - Do not use after the beyond-use date.	☐ DOSING REMINDER - Allow medication to thaw before use. - Use no more than one injection in 24 hours. - Follow your prescriber's dosing instructions.	☑ TELL YOUR HEALTH CARE TEAM Tell your doctor or pharmacist about all medications, over-the-counter products, supplements, and herbal preparations you are taking.

Important: This information is intended for adult patients prescribed Trimix by their physician. It is not intended to cover all possible uses, precautions, drug interactions, or adverse effects. If you have any questions or concerns, contact your doctor or pharmacist.

This is a compounded medication, prescribed just for you. This information has not been evaluated by the Food and Drug Administration.